

**Travis G. Maak, MD
Sports Medicine
University of Utah Orthopaedics
590 Wakara Way
Salt Lake City, UT 84108**

Tel: 801.587.7109 Fax: 801.587.7112

**Hip Arthroscopy Rehabilitation
Capsular Shift with or without FAI –Labral Components**

General Guidelines:

- No external rotation greater than 30 degrees for 4 weeks
- No hyperextension for 4 weeks
- Normalize gait pattern with brace and crutches
- Weight-bearing 20 pounds
- Keep leg from rotating outward when in bed with boot or pillows
- Continuous Passive Motion Machine
 - 4 hours/day or 2 hours if on stationary bike for 2 bouts of 20-30 minute sessions

Frequency of Physical Therapy:

- Seen post-op Day 1
- Seen 1x/week for first month
- Seen 2x/week for second month
- Seen 2-3x/week for third month
- Seen 2-3x/week for fourth month

Precautions following Hip Arthroscopy/FAI: (Refixation/Osteochondroplasty)

- Weight-bearing will be determined by procedure
- Hip flexors tendonitis
- Trochanteric bursitis
- Synovitis
- Manage scarring around portal sites

Travis G. Maak, MD
Sports Medicine
University of Utah Orthopaedics
590 Wakara Way
Salt Lake City, UT 84108

Tel: 801.587.7109 Fax: 801.587.7112

- Increase range of motion focusing on flexion
 - Limit external rotation and hyperextension

Guidelines:

- **Weeks 0-2**

- NO EXTERNAL ROTATION > 20 degrees
- CPM for 4 hours/day
- Bike for 20-30 minutes/day (can be 2x/day)
- Scar massage
- Hip PROM as tolerated with focus on flexion
 - IR as tolerated
 - ER max 20 deg (looking for firm end feel)
- Supine hip log rolling for internal rotation and modified external rotation
- Progress with ROM
 - Introduce stool rotations (AAROM hip IR/ER max to 20)
- Hip isometrics - NO FLEXION
 - Abduction, adduction, extension, ER
- Pelvic tilts
- Supine bridges
- NMES to quads with SAQ with pelvic tilt
- Quadruped rocking for hip flexion
- Gait training PWB with assistive device to normalize gait
- Modalities

- **Weeks 2-4**

Travis G. Maak, MD
Sports Medicine
University of Utah Orthopaedics
590 Wakara Way
Salt Lake City, UT 84108

Tel: 801.587.7109 Fax: 801.587.7112

- Continue with previous therex
- Progress Weight-bearing (week 2)
 - Week 3-4: wean off crutches (2 → 1 → 0) when gait is normalized
- Progress with hip ROM
 - Bent knee fall outs (week 4)
 - Stool rotations for ER (week 3-4) max 30 degrees
 - Prone hip ER/IR (week 4)
- Glut/piriformis stretch
- Progress core strengthening (avoid hip flexor tendonitis)
- Progress with hip strengthening – isotonics all directions except flexion
 - Start isometric sub max pain free hip flexion(3-4 wks)
- Step downs
- Clam shells → isometric side-lying hip abduction
- Hip Hiking (week 4)
- Begin proprioception/balance training
 - Balance boards, single leg stance
- Bike / Elliptical (week 6)
- Scar massage
- Bilateral Cable column rotations (week 4)
- Treadmill side stepping from level surface holding on → inclines (week 5) with good abductor strength
- Aqua therapy in low end of water if available

Travis G. Maak, MD
Sports Medicine
University of Utah Orthopaedics
590 Wakara Way
Salt Lake City, UT 84108

Tel: 801.587.7109 Fax: 801.587.7112

- **Weeks 4-8**

- Continue with previous therex
- Progress with ROM
 - Standing BAPS rotations
 - External rotation with FABER
 - Hip flexor, glute/piriformis, and It-band Stretching – manual and self
 - Hip flexor stretch on stool to increase hip extension
- Progress strengthening LE
 - Introduce hip flexion isotonic (Be aware of hip flexion tendonitis)
 - Multi-hip machine (open/closed chain) consider closed strength if good abductor strength
 - Leg press (bilateral → unilateral)
 - Isokinetics: knee flexion/extension
- Progress core strengthening (avoid hip flexor tendonitis)
 - Prone/side planks
- Progress with proprioception/balance
 - Bilateral → unilateral → foam → dynadisc
- Progress cable column rotations –unilateral → foam
- Side stepping with theraband
- Hip hiking on Stairmaster

- **Weeks 8-12**

- Progressive hip ROM

Travis G. Maak, MD
Sports Medicine
University of Utah Orthopaedics
590 Wakara Way
Salt Lake City, UT 84108

Tel: 801.587.7109 Fax: 801.587.7112

- Hip Joint mobs with mobilization belt into limited ranges of joint motion
ONLY IF NECESSARY
 - Lateral and inferior with rotation when indicated
 - Prone posterior-anterior glides with rotation when indicated
- Progressive LE and core strengthening
- Endurance activities around the hip
- Dynamic balance activities
- **Weeks 12-16**
 - Progressive LE and core strengthening
 - Treadmill running program
 - Sport specific agility drills and plyometrics
- **3,6,12 months Re-Evaluate (Criteria for discharge)**
 - Hip Outcome Score
 - Pain free or at least a manageable level of discomfort
 - MMT within 10 percent of uninvolved LE
 - Biodex test of Quadriceps and Hamstrings peak torque within 15 percent of uninvolved
 - Single leg cross-over triple hop for distance:
 - Score of less than 85% are considered abnormal for male and female
 - Step down test